



## Community Infection Prevention and Control Policy for Care Home settings

# Enteral tube feeding

ENTERAL TUBE FEEDING

**Version 3.00**  
**October 2020**

Please note that the internet version is the only version that is maintained. Any printed copies should, therefore, be viewed as 'uncontrolled' and as such, may not necessarily contain the latest updates and amendments.

**This Policy has been adopted by:**

Organisation: .....

Signed: .....

Job Title: .....

Date Adopted: .....

Review Date: .....

<p>If your organisation would like to exclude or include any additional points to this document, please include below. Please note, the Community IPC team cannot endorse or be held responsible for any addendums.</p>
<p>Version 3.00: Sections 2 and 4.4 amended October 2020.</p>

Community Infection Prevention and Control  
Harrogate and District NHS Foundation Trust  
Gibraltar House, Thurston Road  
Northallerton, North Yorkshire. DL6 2NA  
Tel: 01423 557340  
email: [ipccommunity@hdfnhs.uk](mailto:ipccommunity@hdfnhs.uk)  
[www.infectionpreventioncontrol.co.uk](http://www.infectionpreventioncontrol.co.uk)

**Legal disclaimer**  
This Policy produced by Harrogate and District NHS Foundation Trust is provided 'as is', without any representation endorsement made and without warranty of any kind whether express or implied, including but not limited to the implied warranties of satisfactory quality, fitness for a particular purpose, non-infringement, compatibility, security and accuracy.

These terms and conditions shall be governed by and construed in accordance with the laws of England and Wales. Any dispute arising under these terms and conditions shall be subject to the exclusive jurisdiction of the courts of England and Wales.

## Contents

## Page

1. Introduction.....	4
2. Administration of feeds (aseptic technique).....	4
3. Cleaning and storage of enteral feeding equipment .....	5
4. Administering medication through an enteral feeding tube (clean technique) .....	6
5. Care of the tube insertion site .....	7
6. Infection Prevention and Control resources, education and training.....	8
7. References .....	8

# ENTERAL TUBE FEEDING

## ENTERAL TUBE FEEDING

### 1. Introduction

Enteral tube feeding is a process where nutrition is delivered into an individual's gastrointestinal tract by 1 of 3 ways:

- Through the nose into the stomach by nasogastric feeding tube
- Directly into the stomach by gastrostomy or PEG (percutaneous endoscopic gastrostomy) feeding tube
- Directly into the small bowel by jejunostomy feeding tube

An individualised care plan should be in place for each resident receiving enteral tube feeding.

Only commercially prepared feeds should be used.

Residents receiving enteral tube feeding should be supported by the multi-disciplinary team (MDT).

Enteral tube feeding administration must be undertaken only by suitably trained and competent staff. Initial training and competency should be assessed and monitored by the relevant MDT members.

This Policy for safe practice will assist staff to reduce the risk of infection associated with enteral tube feeding.

It is recommended that regular audits are undertaken. An audit tool is available to download at [www.infectionpreventioncontrol.co.uk](http://www.infectionpreventioncontrol.co.uk).

### 2. Administration of feeds (aseptic technique)

Explain procedure to resident.

1. Check the expiry dates on items to be used, e.g. giving set, feed, (60 ml ENfit reusable enteral syringes, coloured purple, are an example of enteral syringes used in community settings).
2. Ensure appropriate resident positioning, i.e. upper body positioned at a minimum angle of 30° prior to and throughout the feeding period.
3. Take the equipment to the resident's bedside or appropriate private space.
4. Hands must be washed with liquid soap and warm running water and thoroughly dried with paper towels. Alcohol handrub can be used if hands are visibly clean.

5. Put on disposable apron and gloves.
6. Remove the giving set packaging and close the clamp.
7. Aseptic technique to be used, to connect the sterile giving set to the sterile feed container and the enteral feeding tube.
9. Shake the bag/bottle, twist off the cap and without touching the spike, tightly screw on the giving set, which will break the foil seal.
10. Hang the bag on the drip stand, feed the giving set into the pump
11. Prime the giving set, making sure there are no air bubbles.
12. Set the rate of administration as directed by the dietitian
13. Before commencing a feed, slowly flush the feeding tube with a minimum of 30 mls of water - freshly drawn tap water for residents who are not immunosuppressed or sterile water from a freshly opened container for residents who are immunosuppressed. Note: If sterile water is used, it should be labelled with the resident's name, the time and date opened and any unused sterile water should be disposed of after 24 hours.
14. Remove the end cap from the giving set, connect the giving set to the feeding tube and press start.
15. At the end of a feed, slowly flush the feeding tube again with a minimum of 30 mls of water. Tubing should be flushed between intermittent feeds.
16. Feed containers and giving sets are single-use items and should be disposed of at the end of the feeding session.
17. Remove and dispose of gloves, clean hands, remove and dispose of apron, clean hands.
18. After each feed, record the amount of feed and number of flushes given on the resident's prescribing chart.
19. Ensure the resident is comfortable, observe for signs of feeding intolerance.
20. Maintain resident's upper body positioned at a minimum angle of 30° for 1 hour after feeding. Ensure they do not lay flat following feeding period.

### 3. Cleaning and storage of enteral feeding equipment

Feed containers and giving sets are single use items and should be disposed of at the end of the feeding session.

Single use enteral syringes should be disposed of after use.

Single patient use enteral syringes should be:

- Clearly marked with the resident's name, dated at first use and documented accordingly

- Cleaned after each use (as per manufacturer's instructions) – cleaning should be carried out in a suitable sink, which is not used for personal care or routine hand hygiene
- Stored in a clean dry container, marked with the resident's name (the container should be washed with pH neutral detergent and warm water daily)
- Discarded after 1 weeks use or as per manufacturers guidance

The pump used to administer feeds should be cleaned regularly and thoroughly, with pH neutral detergent, e.g. Hospec, and warm water or a detergent wipe.

#### 4. Administering medication through an enteral feeding tube (clean technique)

Administration of drugs via an enteral feeding tube is unlicensed in the UK. If the resident is able to swallow their medication orally in the normal manner, this is preferred. Prescribers should consider alternative routes of administration for which licensed products are available.

Prior to administering medication, check:

- If the resident can take medication orally, if the medication is necessary or if it can be temporarily suspended
- Consider if an alternative route can be used, e.g. buccal (placing between the gums and the cheek), transdermal, topical, rectal or subcutaneous

Medication given via enteral feeding tubes should be administered in liquid or syrup form, suitably diluted, whenever possible. The tube must be flushed thoroughly before and after the administration of medicines.

In the absence of such products, some tablets may be crushed or alternative formulations, such as dispersible tablets, may be considered. Enteric coated or slow release preparations are not suitable for administration via feeding tubes, contact your local pharmacist for further advice.

Some medicines should never be crushed - including modified release tablets, enteric coated tablets, cytotoxic medicines. Prior to preparation, check with a Pharmacist if the medicine is able to be crushed or not.

Where there is a contraindication for medicine to be taken with feed, the dietitian should be consulted to prescribe a suitable regime to ensure that the resident's nutritional requirements are maintained.

To avoid interaction between medicines and feed, do not add medication directly into the enteral feed.

Administering medication (clean technique):

1. Hands should be washed thoroughly with liquid soap and warm running water or an alcohol handrub used before administering medication
2. Put on gloves and, if there is a risk of contamination to the uniform, apron
3. If an enteral feed is in progress, stop the feed. If the administration of any of the medicine with feed is contraindicated, ensure the suggested time of withholding the feed is adhered to.
4. Using an enteral syringe, slowly flush the feeding tube with a minimum of 30 mls of water - freshly drawn tap water for residents who are not immunosuppressed or either cooled freshly boiled water or sterile water from a freshly opened container for residents who are immunosuppressed  
Note: If sterile water is used, it should be labelled with the resident's name, the time and date opened and any unused sterile water should be disposed of after 24 hours.
5. Prepare each medication to be given separately to avoid interaction of the drugs and ensure solubility:
  - Soluble tablets - dissolve in 10-15 mls water
  - Tablets - crush with a clean mortar and pestle or tablet crusher
  - Liquids - shake well. Mix thick liquids with an equal volume of water
6. Administer the medication through the feeding tube, using a 50 ml syringe
7. A 10 ml flush of water should be given in between each medication to prevent tube blockages and drug interactions
8. Flush the feeding tube with 30 mls of water at the end of administering the medications
9. Remove and dispose of gloves, clean hands, remove and dispose of apron, clean hands
10. Record on resident's prescribing chart

## 5. Care of the tube insertion site

Hand hygiene is essential, before contact with the resident's enteral feeding tube and/or insertion site. Hands should be washed with liquid soap and warm running water and thoroughly dried using paper towels. Alcohol handrub can be used if hands are visibly clean.

If there is pain on feeding or external leakage of stomach contents, or fresh bleeding, stop any feed immediately and urgently contact your local Hospital Emergency Department.

Unless advised differently by the enteral tube manufacturers or a healthcare professional, the alternative advice should be documented in the resident's care plan:

- Following insertion of an enteral tube, treat the insertion site as a surgical wound using an aseptic technique for the first 48 hours, keeping it clean and dry
- After 48 hours, the insertion site should be washed daily with tap water and dried thoroughly. Avoid water immersion, e.g. bathing in a bath, for 2 weeks and reapply dressing
- When the insertion site has healed, usually 10-12 days, no dressing is required. The site should be inspected and cleaned with soap and water daily
- Weekly 360° rotation of the tube (and in some cases 2-4 cm advancement of the tube prior to rotation), is recommended to prevent adherence, and to avoid infection related to 'buried bumper syndrome'

## 6. Infection Prevention and Control resources, education and training

The Community Infection Prevention and Control (IPC) Team have produced a wide range of innovative educational and IPC resources designed to assist your Care Home in achieving compliance with *The Health and Social Care Act 2008: Code of Practice on the prevention and control of infections and related guidance* and CQC registration requirements.

These resources are either free to download from the website or available at a minimal cost covering administration and printing:

- 29 IPC Policy documents for Care Home settings
- 'Preventing Infection Workbook: Guidance for Care Homes'
- 'IPC CQC Inspection Preparation Pack for Care Homes'
- IPC audit tools, posters, leaflets and factsheets
- 'IPC Bulletin for Care Homes'

In addition, we hold educational study events in North Yorkshire and can arrange bespoke training packages and 'Mock IPC CQC Inspections'. Prices vary depending on your requirements and location.

Further information on these high quality evidence-based resources is available at [www.infectionpreventioncontrol.co.uk](http://www.infectionpreventioncontrol.co.uk).

## 7. References

Department of Health (2015) *The Health and Social Care Act 2008: Code of Practice on the prevention and control of infections and related guidance*



Department of Health and Health Protection Agency (2013) *Prevention and control of infection in care homes – an information resource*

MGP Ltd (January 2019) *Medication management of patients with nasogastric (NG), percutaneous endoscopic gastrostomy (PEG), or other enteral feeding tubes*

National Institute for Health and Care Excellence (2012, updated February 2017) *Healthcare-associated Infections: prevention and control in primary and community care CG139*

National Institute for Health and Care Excellence (2006, updated August 2017) *Nutrition support for adults: oral nutrition support, enteral tube feeding and parental nutrition CG32*

National Patient Safety Agency (March 2010) *NPSA/2010/RRR010 Rapid Response Report: Early detection of complications after gastrostomy*

Royal Marsden NHS Foundation Trust (2020) *The Royal Marsden Hospital Manual of Clinical and Cancer Nursing Procedures 10<sup>th</sup> Edition*