



Community Infection Prevention and Control Policy for General Practice

(also suitable for adoption by other healthcare providers, e.g. Dental Practice, Podiatry)

Venepuncture

Version 2.00 July 2021 Please note that the internet version is the only version that is maintained. Any printed copies should, therefore, be viewed as 'uncontrolled' and as such, may not necessarily contain the latest updates and amendments.

This document has been adopted as a Policy by: Organisation: Job Title: Date Adopted: Review Date: If your organisation would like to exclude or include any additional points to this document, please include below. Please note, the Community IPC Team cannot endorse or be held responsible for any addendums.

Community Infection Prevention and Control Harrogate and District NHS Foundation Trust Gibraltar House, Thurston Road Northallerton, North Yorkshire, DL6 2NA

Tel: 01423 557340

email: infectionprevention.control@nhs.net www.infectionpreventioncontrol.co.uk

Legal Disclaimer

This Policy produced by Harrogate and District NHS Foundation Trust is provided 'as is', without any representation endorsement made and without warranty of any kind whether express or implied, including but not limited to the implied warranties of satisfactory quality, fitness for a particular purpose, noninfringement, compatibility, security and accuracy.

These terms and conditions shall be governed by and construed in accordance with the laws of England and Wales. Any dispute arising under these terms and conditions shall be subject to the exclusive jurisdiction of the courts of England and Wales.

Contents Page

1.	Introduction	4
	Gloves	
3.	Skin decontamination	4
4.	Skin dressing	5
5.	Preparation for venepuncture	5
6.	Procedure	5
7.	Infection Prevention and Control resources, education and training	7
8.	References	7

VENEPUNCTURE

1. Introduction

Venepuncture is the procedure of entering a vein with a needle in order to obtain a sample of blood for diagnostic purposes.

Venepuncture breaches the circulatory system, therefore, to minimise the risk of injury and/or infection to both staff and patients, standard infection control precautions and transmission based precautions (SICPs and TBPs) should be adhered to.

The procedure should only be undertaken by appropriately trained and competent staff.

Reusable tourniquets can harbour microorganisms, such as bacteria and viruses, and, therefore pose a risk of transmission of infection. Practices should risk assess the use of reusable tourniquets against single use ones, and if reusable tourniquets are used, a schedule for their decontamination and replacement should be implemented.

2. Gloves

Best practice is to wear non-sterile disposable gloves, nitrile or latex, when undertaking venepuncture, refer to the 'PPE Policy for General Practice'. The wearing of gloves has been shown to reduce the volume of blood transferred in a needlestick injury by 52% compared with not wearing gloves, which can help reduce the risk of acquiring a blood-borne virus (BBV) if you sustain a needlestick injury.

3. Skin decontamination

Best practice is to clean the skin:

- Good skin preparation reduces the risk of infection from colonising skin bacteria
- Use a 70% isopropyl alcohol or 2% chlorhexidine in 70% alcohol skin wipe to clean the area for 30 seconds using moderate pressure and allow to air dry

4. Skin dressing

Applying a lint free sterile dressing over the puncture site is important to prevent infection.

5. Preparation for venepuncture

Ensure all equipment is available:

- A single use or clean reusable tourniquet
- Vacuum blood collection system needles, preferably a safer sharps device to reduce the risk of accidental sharps injury, and appropriate blood bottles
- · Clean tray or receiver
- Sharps container
- Non-sterile disposable latex or nitrile gloves
- Nitrile gloves should be available for those with a confirmed or suspected latex allergy
- Disposable apron
- 70% isopropyl alcohol or 2% chlorhexidine in 70% alcohol skin wipes
- Sterile gauze
- Sterile adhesive plaster or hypoallergenic tape

Check the patient's details and discuss the procedure. Check for any previous problems with venepuncture, any allergies and obtain verbal consent.

Note: Blood must be drawn in a specific order, 'the order of draw', to avoid cross-contamination of additives from one tube to another, affecting laboratory results. Follow your blood collection systems guidance for order of draw.

6. Procedure

- Assemble specific equipment required, checking packaging for damage and expiry dates.
- Wash hands with liquid soap and warm running water or use alcohol handrub, using the technique given in the 'Hand hygiene Policy for General Practice'.
- Put on disposable apron.

- Position the patient comfortably and support the limb. If a pillow is used, it should be free from a linen pillow case and have a wipeable cover.
- Apply tourniquet, 5-10 cm above the proposed puncture site.
- Palpate the proposed puncture site to select a vein.
- Wash hands or use alcohol handrub.
- Put on disposable gloves.
- Clean patient's skin with 70% isopropyl alcohol or 2% chlorhexidine in 70% alcohol skin wipes, using a moderate pressure rubbing action (dependent upon skin assessment/fragility) for 30 seconds and allow the skin to dry.
- Do not re-palpate or touch the area after cleaning.
- Remove cover from needle or needle guard and inspect the device for any faults.
- Anchor the vein by applying manual traction on the skin a few centimetres below the proposed puncture site.
- Insert needle smoothly and undertake the procedure of venepuncture as per training received and vacuum system being used.
- Release the tourniquet (if not previously done).
- Place a low linting sterile gauze over puncture site.
- Remove the needle, but do not apply pressure until fully removed.
- Discard sharps in a sharps container as per 'Safe management of sharps and inoculation injuries Policy for General Practice'.
- Apply pressure and use additional piece of sterile gauze if necessary.
 Discourage patient from bending their arm.
- Gently invert blood bottle tube at least six times.
- Inspect puncture point for bleeding.
- Apply sterile adhesive plaster or hypoallergenic tape.
- · Remove and dispose of gloves and apron.
- Wash hands with liquid soap and warm running water or use alcohol handrub.
- Label sample bottles.
- Complete the specimen request form.

7. Infection Prevention and Control resources, education and training

The Community Infection Prevention and Control (IPC) Team have produced a wide range of innovative educational and IPC resources designed to assist your Practice in achieving compliance with *The Health and Social Care Act 2008: Code of Practice on the prevention and control of infection and related guidance* and CQC registration requirements.

These resources are either free to download from the website or available at a minimal cost covering administration and printing:

- 25 IPC Policy documents for General Practice
- 'Preventing Infection Workbook: Guidance for General Practice'
- 'IPC CQC inspection preparation Pack for General Practice'
- IPC audit tools, posters, leaflets and factsheets
- 'IPC Bulletin for GP Practice Staff'

In addition, we hold educational study events in North Yorkshire and York and can arrange bespoke training packages and 'Mock IPC CQC Inspections'. Prices vary depending on your requirements and location.

Further information on these high quality evidence-based resources is available at www.infectionpreventioncontrol.co.uk.

8. References

Hospital Infection Society (2007) Standardization of needlestick injury and evaluation of a novel virus-inhibiting protective glove *Journal of Hospital Infection Volume 66, Issue 4, August 2007, {Pages 339-345)*

Loveday HP, et al, epic 3: National Evidence-Based Guidelines for Preventing Healthcare-Associated Infections in NHS Hospitals in England Journal of Hospital Infection 86S1 (2014) S1-S70

Royal Marsden NHS Foundation Trust (2020) *The Royal Marsden Hospital Manual of Clinical and Cancer Nursing Procedures* 10th Edition