



Community Infection Prevention and Control Policy for General Practice

(also suitable for adoption by other healthcare providers, e.g. Dental Practice, Podiatry)

Respiratory illnesses

Version 1.00 November 2023 Please note that the internet version is the only version that is maintained. Any printed copies should, therefore, be viewed as 'uncontrolled' and as such, may not necessarily contain the latest updates and amendments.

This Policy has been adopted b	y:
Organisation:	
Signature:	Name:
Job title:	
Adoption date:	
Review date:	
through local governance channel Infection Prevention and Control (be risk assessed, evidence-based, authorised is and documented below. The Community IPC) Team, Harrogate and District NHS or be held responsible for any variations.

Community Infection Prevention and Control Harrogate and District NHS Foundation Trust Gibraltar House, Thurston Road Northallerton, North Yorkshire. DL6 2NA Tel: 01423 557340

email: infectionprevention.control@nhs.net www.infectionpreventioncontrol.co.uk

Legal disclaimer

This Policy produced by Harrogate and District NHS Foundation Trust is provided 'as is', without any representation endorsement made and without warranty of any kind whether express or implied, including but not limited to the implied warranties of satisfactory quality, fitness for a particular purpose, non-infringement, compatibility, security and accuracy.

These terms and conditions shall be governed by and construed in accordance with the laws of England and Wales. Any dispute arising under these terms and conditions shall be subject to the exclusive jurisdiction of the courts of England and Wales.

Contents Page

1.	Introduction	4
2.	What are respiratory tract infections?	4
3.	How are respiratory illnesses spread?	5
4.	Management of a patient	5
5.	Good respiratory and cough hygiene	6
6.	TBPs for respiratory illnesses	6
7.	Infection Prevention and Control resources, education and training	6
8.	References	7
9.	Appendices	7
Арре	endix 1: A-Z respiratory illnesses	. 8

RESPIRATORY ILLNESSES

1. Introduction

Respiratory illnesses are a major cause of hospitalisation, morbidity and death in the elderly and are amongst the most common winter ailments. Ranging from a self-limiting, mild, coryzal like illness such as rhinovirus (common cold) to severe bronchitis, bronchiolitis and pneumonia.

Those suffering from underlying chronic health conditions become more susceptible and vulnerable to severe disease. Vaccination provides the best protection against acquiring and spreading infection.

'Standard infection control precautions' (SICPs), especially 'Respiratory and cough hygiene' alongside good ventilation, will help to reduce the risk of spreading respiratory illnesses from an infected person to others.

SICPs may, however be insufficient to prevent transmission of specific infections and additional 'Transmission based precautions' (TBPs) may be required. Refer to the 'SICPs and TBPs Policy for General Practice'.

When caring for patients in relation to any new or emerging infection, staff should refer to the latest national infection prevention and control guidance.

2. What are respiratory tract infections?

Respiratory tract infections (RTIs) are infections of parts of the body involved in breathing, such as the sinuses, throat, airways or lungs. RTIs are mainly caused by viruses and can affect the upper respiratory tract or the lower respiratory tract.

Upper respiratory tract infections

Upper respiratory tract infections (URTIs) include the common cold, tonsillitis, sinusitis, laryngitis and flu.

The most common symptoms are headache, aching muscles, a blocked/stuffed up or runny nose, sneezing and a sore throat. URTIs caused by a virus, e.g. the common cold, usually get better without any treatment.

Lower respiratory tract infections

Lower respiratory tract infections (LRTIs) include bronchitis (an infection of the airways), pneumonia (lung infection), bronchiolitis (an infection of the small

airways that affects babies and children) and tuberculosis, a bacterial lung infection. Flu can affect both the upper and lower respiratory tract.

The most common symptom of LRTI is coughing, in severe cases patients cough up mucus and can suffer from breathlessness, wheezing and chest tightness.

RTIs caused by bacteria, e.g. pneumonia, tuberculosis, often require antibiotic treatment and in some cases, admission to hospital. Bacterial infections are not covered in this policy.

3. How are respiratory illnesses spread?

Respiratory illnesses are spread by:

- Predominantly droplet transmission. Droplets are generated during coughing, sneezing, talking. If droplets from an infected person come into contact with the mucous membranes, e.g. eyes, nose, mouth, of another person, they can cause infection. Droplets remain in the air for a short period and can travel approximately 1 metre. They can land on surfaces and equipment and if touched, infect others if that person has contact with their eyes, nose, mouth
- Aerosol transmission is usually associated with an aerosol generating procedure (AGP). An AGP can result in the release of airborne particles (aerosols) from the respiratory tract, when treating someone with a confirmed or suspected viral infection. During an AGP, smaller viral particles than droplets are produced which can remain suspended in the air for longer and travel further than 1 metre

Spirometry, taking a diagnostic throat/nose swab, the use of nebulisers and similar procedures carried out in a GP Practice, are not usually categorised as AGPs.

4. Management of a patient

To help reduce the spread of respiratory illnesses, 'Standard infection control precautions' (SICPs) and, where required, 'Transmission based precautions' (TBPs) should always be followed.

- Staff must wear appropriate personal protective equipment (PPE), as per national guidance, including a fluid resistant surgical mask (FRSM) and eye protection, e.g. goggles or visor (prescription glasses do not provide adequate protection).
- Advise the patient to wear a face mask and if possible keep it on throughout the visit.
- Place the patient in a separate waiting room, consulting room or area away

from others.

- Ensure good ventilation of the room by opening windows.
- After the patient leaves the room, keep the window open, clean and disinfect
 the immediate area and any reusable devices, care equipment, treatment
 couch, etc.
- If transfer to hospital is required, the ambulance service and hospital department should be informed of any respiratory illness.

For advice on influenza management, please contact your local Community Infection Prevention and Control (IPC) or UK Health Security Agency (UKHSA) Team.

Other illnesses caused by viruses, e.g. chicken pox, measles, are also spread by inhaling respiratory secretion droplets from an infected person's cough or sneeze, or from touching surfaces contaminated when they coughed or sneezed.

Good respiratory and cough hygiene

Please refer to the 'Respiratory and cough hygiene Policy for GP Practice'.

6. TBPs for respiratory illnesses

Appendix 1 provides an A-Z listing of the common respiratory virus infections. It specifies the type of precautions, optimal patient placement and PPE required.

Refer to the 'SICPs and TBPs Policy for GP Practice'.

7. Infection Prevention and Control resources, education and training

The Community IPC Team have produced a wide range of innovative educational and IPC resources designed to assist your General Practice in achieving compliance with the *Health and Social Care Act 2008: code of practice on the prevention and control of infection and related guidance* and CQC registration requirements.

These resources are either free to download from the website or available at a minimal cost covering administration and printing:

27 IPC Policy documents for General Practice

- Preventing Infection Workbook: Guidance for General Practice
- IPC CQC inspection preparation Pack for General Practice
- IPC audit tools, posters, leaflets and factsheets
- IPC Bulletin for General Practice Staff

In addition, we hold educational study events in North Yorkshire.

Further information on these high quality evidence-based resources is available at www.infectionpreventioncontrol.co.uk.

8. References

Department of Health and Social Care (Updated December 2022) Health and Social Care Act 2008: code of practice on the prevention and control of infections and related guidance

NHS England (2022, updated 2023) National infection prevention and control manual (NIPCM) for England

NHS Respiratory tract infections (RTIs) <u>www.nhs.uk/conditions/Respiratory-tract-infection/</u>

9. Appendices

Appendix 1: A-Z respiratory illnesses

Appendix 1: A – Z respiratory illnesses								
Virus/Agent	Incubation	Required	Duration (very old/ immunocompromised patients will shed the virus for longer)	PPE				
Coronavirus (non- COVID), enterovirus, bocavirus	3-5 days	No	No	SICPs				
COVID-19	1-14 days	Droplet/Airborne	Up to 10 days from onset of symptoms	Gloves, apron, eye protection and FRSM for routine care. FFP3 for AGPs*				
Human metapneumovirus	4-6 days	No	No	SICPs				
Influenza A/B (flu)	1-3 days	Droplet	5 days from onset of symptoms if on antiviral treatment. 7 days from onset of symptoms if not on antiviral treatment. Immunocompromised patients require 10 days isolation	Gloves, apron, eye protection and FRSM for routine care. FFP3 for AGPs*				
Parainfluenza (types 1-4)	1-7 days	Droplet	5 days	Gloves, apron, eye protection and FRSM for routine care. FFP3 for AGPs*				
Respiratory syncytial virus RSV)	3-5 days	Droplet	5 days	Gloves, apron, eye protection and FRSM for routine care. FFP3 for AGPs*				
Rhinovirus/adenovirus (common cold)	1-2 days	No	No	SICPs				

^{*}AGPs (aerosol generating procedures) are rare in a GP Practice, but include respiratory suctioning beyond the oropharynx and tracheostomy tube insertion/removal.